

STATE OF OHIO IMMUNIZATION EXEMPTION

Per OHIO STATUTE 3313.671

Religious, Reasons of Conscience, and Medical Exemption Form

(4) A pupil who presents a written statement of the pupil's parent or guardian in which the parent or guardian declines to have the pupil immunized or reasons of conscience, including religious convictions, is not required to be immunized.

(5) A child whose physician certifies in writing that such immunization against any disease is medically contradicted is not required to be immunized against that disease.

I understand that the immunization Law permits me to sign a waiver on my child taking the immunization.

I hereby object and request the school to waiver the immunization of my child against the following:

| | | | | |
|--------|---------------|------------------------|-------------|--------------|
| D.P.T. | Polio | Hepatitis A | MMR | Pneumococcal |
| Dtap | Meningococcal | Varicella (chickenpox) | Flu Vaccine | Rotavirus |

Child's Name: _____

MUST SELECT AT LEAST ONE REASON:

- RELIGIOUS – List name of denomination _____
- REASON OF CONSCIENCE – Please explain _____

- MEDICAL REASON – You must have a signed statement from your physician stating the condition and attach it to this form.

I further understand that during the course of a an outbreak of any of the aforementioned vaccine preventable diseases, that the student named here is subject to exclusion from school for the duration of the outbreak.

This action is necessary not only to protect this student, but the remainder of the students and faculty of the school.

Parent/Guardian Signature _____ Date: _____

Address: _____

Medication Administration Record (MAR)
General Medication Form
(Including Asthma Inhaler and Epinephrine Autoinjector Use)

Student Information

| | | | |
|---|-------------|---------|---------------|
| Student name | | | Date of birth |
| Student address | | | |
| School | Grade/Class | Teacher | School year |
| List any known drug allergies/reactions | | Height | Weight |

Prescriber Authorization

| | | | |
|---|--|---------------|-----|
| Name of medication | Circumstance for use | | |
| Dosage | Route | Time/Interval | |
| Date to begin medication | Date to end medication | | |
| Circumstances for use | | | |
| Special instructions | | | |
| Treatment in the event of an adverse reaction | | | |
| Epinephrine Autoinjector | <input type="checkbox"/> Not applicable <input type="checkbox"/> Yes, as the prescriber I have determined that this student is capable of possessing and using this autoinjector appropriately and have provided the student with training in the proper use of the autoinjector. | | |
| Asthma Inhaler | <input type="checkbox"/> Not applicable <input type="checkbox"/> Yes, if conditions are satisfied per ORC 3317.716, the student may possess and use the inhaler at school or at any activity event or program sponsored by or in which the student's school is a participant. | | |
| Procedures for school employees if the student is unable to administer the medication or if it does not produce the expected relief | | | |
| Possible Severe Adverse Reaction(s) per ORC 3317.716 and 3313.718 | | | |
| a) To the student for whom it is prescribed (that should be reported to the prescriber) | | | |
| b) To a student for whom it is not prescribed who receives a dose | | | |
| Other medication instructions | | | |
| Does medication require refrigeration? <input type="checkbox"/> Yes <input type="checkbox"/> No Is the medication a controlled substance? <input type="checkbox"/> Yes <input type="checkbox"/> No | | | |
| Prescriber signature | Date | Phone | Fax |
| Prescriber name (print) | | | |
| Reminder note for prescriber: ORC 3313.718 requires backup epinephrine autoinjector and best practice recommends backup asthma inhaler. | | | |

Parent/Guardian Authorization

| | | | |
|--|------|------------------|------------------|
| <input checked="" type="checkbox"/> I authorize an employee of the school board to administer the above medication. <input checked="" type="checkbox"/> I understand that additional parent/prescriber signed statements will be necessary if the dosage of medication is changed. <input checked="" type="checkbox"/> I also authorize the licensed healthcare professional to talk with the prescriber or pharmacist to clarify medication order. | | | |
| <input checked="" type="checkbox"/> Medication form must be received by the principal, his/her designee, and/or the school nurse. <input checked="" type="checkbox"/> I understand that the medication must be in the original container and be properly labeled with the student's name, prescriber's name, date of prescription, name of medication, dosage, strength, time interval, route of administration and the date of drug expiration when appropriate. | | | |
| Parent/Guardian signature | Date | #1 contact phone | #2 contact phone |

Parent/Guardian Self-Carry Authorization

| | | | |
|---|------|------------------|------------------|
| <input type="checkbox"/> For Epinephrine Autoinjector: As the parent/guardian of this student, I authorize my child to possess and use an epinephrine autoinjector, as prescribed, at the school and any activity, event, or program sponsored by or in which the student's school is a participant. I understand that a school employee will immediately request assistance from an emergency medical service provider if this medication is administered. I will provide a backup dose of the medication to the school principal or nurse as required by law. | | | |
| <input type="checkbox"/> For Asthma Inhaler: As the parent/guardian of this student, I authorize my child to possess and use an asthma inhaler as prescribed, at the school and any activity, event, or program sponsored by or in which the student's school is a participant. | | | |
| Parent/Guardian signature | Date | #1 contact phone | #2 contact phone |


ASTHMA ACTION PLAN



Asthma and Allergy
Foundation of America
aafa.org

| | |
|-----------------------|-------------------|
| Name: | Date: |
| Doctor: | Medical Record #: |
| Doctor's Phone #: Day | Night/Weekend |
| Emergency Contact: | |
| Doctor's Signature: | |

The colors of a traffic light will help you use your asthma medicines.



GREEN means Go Zone!
Use preventive medicine.

YELLOW means Caution Zone!
Add quick-relief medicine.

RED means Danger Zone!
Get help from a doctor.

Personal Best Peak Flow: _____

GO Use these daily controller medicines:

You have *all* of these:

- Breathing is good
- No cough or wheeze
- Sleep through the night
- Can work & play

Peak flow:

from _____
to _____

| MEDICINE | HOW MUCH | HOW OFTEN/WHEN |
|---------------------------------|----------|----------------|
| | | |
| | | |
| | | |
| For asthma with exercise, take: | | |
| | | |

CAUTION Continue with green zone medicine and add:

You have *any* of these:

- First signs of a cold
- Exposure to known trigger
- Cough
- Mild wheeze
- Tight chest
- Coughing at night

Peak flow:

from _____
to _____

| MEDICINE | HOW MUCH | HOW OFTEN/ WHEN |
|---------------------------------|----------|-----------------|
| | | |
| | | |
| | | |
| CALL YOUR ASTHMA CARE PROVIDER. | | |

DANGER Take these medicines and call your doctor now.

Your asthma is getting worse fast:

- Medicine is not helping
- Breathing is hard & fast
- Nose opens wide
- Trouble speaking
- Ribs show (in children)

Peak flow:

reading below _____

| MEDICINE | HOW MUCH | HOW OFTEN/WHEN |
|----------|----------|----------------|
| | | |
| | | |
| | | |

GET HELP FROM A DOCTOR NOW! Your doctor will want to see you right away. It's important! If you cannot contact your doctor, go directly to the emergency room. DO NOT WAIT.

Make an appointment with your asthma care provider within two days of an ER visit or hospitalization.

**PLACE
PICTURE
HERE**

Name: _____ D.O.B.: _____

Allergic to: _____

Weight: _____ lbs. Asthma: Yes (higher risk for a severe reaction) No

NOTE: Do not depend on antihistamines or inhalers (bronchodilators) to treat a severe reaction. USE EPINEPHRINE.

Extremely reactive to the following allergens: _____

THEREFORE:

- If checked, give epinephrine immediately if the allergen was **LIKELY** eaten, for **ANY** symptoms.
- If checked, give epinephrine immediately if the allergen was **DEFINITELY** eaten, even if no symptoms are apparent.

FOR ANY OF THE FOLLOWING:
SEVERE SYMPTOMS



LUNG

Shortness of breath, wheezing, repetitive cough



HEART

Pale or bluish skin, faintness, weak pulse, dizziness



THROAT

Tight or hoarse throat, trouble breathing or swallowing



MOUTH

Significant swelling of the tongue or lips



SKIN

Many hives over body, widespread redness



GUT

Repetitive vomiting, severe diarrhea



OTHER

Feeling something bad is about to happen, anxiety, confusion

**OR A
COMBINATION**
of symptoms
from different
body areas.



1. **INJECT EPINEPHRINE IMMEDIATELY.**
2. **Call 911.** Tell emergency dispatcher the person is having anaphylaxis and may need epinephrine when emergency responders arrive.
 - Consider giving additional medications following epinephrine:
 - » Antihistamine
 - » Inhaler (bronchodilator) if wheezing
 - Lay the person flat, raise legs and keep warm. If breathing is difficult or they are vomiting, let them sit up or lie on their side.
 - If symptoms do not improve, or symptoms return, more doses of epinephrine can be given about 5 minutes or more after the last dose.
 - Alert emergency contacts.
 - Transport patient to ER, even if symptoms resolve. Patient should remain in ER for at least 4 hours because symptoms may return.

MILD SYMPTOMS



NOSE

Itchy or runny nose, sneezing



MOUTH

Itchy mouth



SKIN

A few hives, mild itch



GUT

Mild nausea or discomfort

**FOR MILD SYMPTOMS FROM MORE THAN ONE
SYSTEM AREA, GIVE EPINEPHRINE.**

**FOR MILD SYMPTOMS FROM A SINGLE SYSTEM
AREA, FOLLOW THE DIRECTIONS BELOW:**

1. Antihistamines may be given, if ordered by a healthcare provider.
2. Stay with the person; alert emergency contacts.
3. Watch closely for changes. If symptoms worsen, give epinephrine.

MEDICATIONS/DOSES

Epinephrine Brand or Generic: _____

Epinephrine Dose: 0.1 mg IM 0.15 mg IM 0.3 mg IM

Antihistamine Brand or Generic: _____


Antihistamine Dose: _____

Other (e.g., inhaler-bronchodilator if wheezing): _____




HOW TO USE AUVI-Q® (EPINEPHRINE INJECTION, USP), KALEO

1. Remove Auvi-Q from the outer case. Pull off red safety guard.
2. Place black end of Auvi-Q against the middle of the outer thigh.
3. Press firmly until you hear a click and hiss sound, and hold in place for 2 seconds.
4. Call 911 and get emergency medical help right away.




HOW TO USE EPIPEN®, EPIPEN JR® (EPINEPHRINE) AUTO-INJECTOR AND EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF EPIPEN®), USP AUTO-INJECTOR, MYLAN AUTO-INJECTOR, MYLAN

1. Remove the EpiPen® or EpiPen Jr® Auto-Injector from the clear carrier tube.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, remove the blue safety release by pulling straight up.
3. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
4. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE IMPAX EPINEPHRINE INJECTION (AUTHORIZED GENERIC OF ADRENALCLICK®), USP AUTO-INJECTOR, AMNEAL PHARMACEUTICALS

1. Remove epinephrine auto-injector from its protective carrying case.
2. Pull off both blue end caps: you will now see a red tip. Grasp the auto-injector in your fist with the red tip pointing downward.
3. Put the red tip against the middle of the outer thigh at a 90-degree angle, perpendicular to the thigh. Press down hard and hold firmly against the thigh for approximately 10 seconds.
4. Remove and massage the area for 10 seconds. Call 911 and get emergency medical help right away.




HOW TO USE TEVA'S GENERIC EPIPEN® (EPINEPHRINE INJECTION, USP) AUTO-INJECTOR, TEVA PHARMACEUTICAL INDUSTRIES

1. Quickly twist the yellow or green cap off of the auto-injector in the direction of the "twist arrow" to remove it.
2. Grasp the auto-injector in your fist with the orange tip (needle end) pointing downward. With your other hand, pull off the blue safety release.
3. Place the orange tip against the middle of the outer thigh at a right angle to the thigh.
4. Swing and push the auto-injector firmly into the middle of the outer thigh until it 'clicks'. Hold firmly in place for 3 seconds (count slowly 1, 2, 3).
5. Remove and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.



HOW TO USE SYMJEPi™ (EPINEPHRINE INJECTION, USP)

1. When ready to inject, pull off cap to expose needle. Do not put finger on top of the device.
2. Hold SYMJEPi by finger grips only and slowly insert the needle into the thigh. SYMJEPi can be injected through clothing if necessary.
3. After needle is in thigh, push the plunger all the way down until it clicks and hold for 2 seconds.
4. Remove the syringe and massage the injection area for 10 seconds. Call 911 and get emergency medical help right away.
5. Once the injection has been administered, using one hand with fingers behind the needle slide safety guard over needle.



ADMINISTRATION AND SAFETY INFORMATION FOR ALL AUTO-INJECTORS:

1. Do not put your thumb, fingers or hand over the tip of the auto-injector or inject into any body part other than mid-outer thigh. In case of accidental injection, go immediately to the nearest emergency room.
2. If administering to a young child, hold their leg firmly in place before and during injection to prevent injuries.
3. Epinephrine can be injected through clothing if needed.
4. Call 911 immediately after injection.

OTHER DIRECTIONS/INFORMATION (may self-carry epinephrine, may self-administer epinephrine, etc.):

Treat the person before calling emergency contacts. The first signs of a reaction can be mild, but symptoms can worsen quickly.

EMERGENCY CONTACTS — CALL 911

RESCUE SQUAD: _____

DOCTOR: _____ PHONE: _____

PARENT/GUARDIAN: _____ PHONE: _____

OTHER EMERGENCY CONTACTS

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____

NAME/RELATIONSHIP: _____ PHONE: _____